

Comment on: Modular Theory and Geometry

K. Ebrahimi-Fard ¹

*Physikalisches Institut der Universität Bonn
Nußallee 12, D-53115 Bonn, Germany*

Abstract

In this note we comment on part of a recent article by B. Schroer and H.-W. Wiesbrock. Therein they calculate some new modular structure for the $U(1)$ -current-algebra (Weyl-algebra). We point out that their findings are true in a more general setting.

¹ e-mail: fard@th.physik.uni-bonn.de

I We would like to add a point to a recent inspiring work of B. Schroer and H.-W. Wiesbrock [1]² implying a modular origin of the chiral "higher dilation" diffeomorphisms.

For the $U(1)$ -current-algebra in two-dimensional spacetime they construct states invariant under higher representations of the Möbius-group, generated by the modes $L_{-n,0,n}$. The new states fulfill the KMS-property with respect to modified dilations. These modified dilations are identified with the modular group associated to von Neumann algebras localized in well-chosen regions and the new states.

In the original version [1] of the conjecture it was overlooked that the chosen state lacked the property of faithfulness if used on doubly-localized intervals. As far as the demonstration of the modular origin of the diffeomorphism group was concerned this was corrected in the work of Schroer and Fassarella [3] but the correction was at the expense of the original conjecture. Here we show that due to the geometrical properties of the modified transformations respectively the new F-S states the use of the split-property [16] allows a faithful extension of these F-S states to doubly-localized intervals and in this way the demonstration of the modular origin of the diffeomorphism group is harmonized with the understanding of the modular structure of double intervals.

In section 2 we sketch briefly the ansatz and result concerning the modular structure in the case of the $U(1)$ -current-algebra. Section 3 contains our point to add, providing a more general point of view of the aforementioned results of Schroer and Wiesbrock. In section 4 we investigate so-called multilocalized algebras by using the split-property. The product states and modular group of such algebras are identified. Some additional remarks and a short summary are given in Section 5.

II Conformal field theory in two dimensions (CFT_2) [15] provides a well suited realm for algebraic quantum field theory [2]³, especially for problems concerning the geometric identification of the modular structure [4, 5]. Minkowskian CFT_2 may be represented on the product of two circles, $S^1 \times S^1$ -spacetime (the "compact- picture"). The global symmetry group of the CFT_2 is the Möbius-group $PSU(1, 1) \times PSU(1, 1)$. We will concentrate on one of the groups, being realized on one of the circles:

$$PSU(1, 1) := SU(1, 1)/\{\pm 1\},$$

$$SU(1, 1) := \left\{ \begin{pmatrix} \alpha & \beta \\ \bar{\beta} & \bar{\alpha} \end{pmatrix} \middle| \alpha, \beta \in \mathbb{C}, |\alpha|^2 - |\beta|^2 = 1 \right\}.$$

² Recently this work was corrected in an important point and also extended by Schroer and Fassarella [3]. Their new results demanded also for a correction of an earlier version of this paper.

³ For a very recent review of the current state of algebraic QFT see [6].

The spectrum generating algebra of reparametrizations of the circle is generated by the Virasoro algebra (with central charge c) $\mathfrak{L}_c$:

$$[L_n, L_m] = (n - m)L_{n+m} + \frac{c}{12}(n^3 - n)\delta_{n+m,0}, \quad n \in \mathbb{Z}. \quad (1)$$

The globally realized group $PSU(1, 1)$ has the underlying generators L_{-1}, L_0, L_1 , fulfilling a $sl(2, \mathbb{C})$ -algebra:

$$[L_1, L_{-1}] = 2L_0, \quad [L_{\pm 1}, L_0] = \pm L_{\pm 1}. \quad (2)$$

The Virasoro algebra $\mathfrak{L}_c$ contains infinitely many further $sl(2, \mathbb{C})$ -algebras, generated by the modes L_{-n}, L_0, L_n , $n > 1$:

$$\begin{aligned} L_{-n} &\mapsto \tilde{L}_{-n} := \frac{1}{n}L_{-n} \\ L_0 &\mapsto \tilde{L}_0 := \frac{1}{n}L_0 + \frac{c}{24}\frac{(n^2-1)}{n} \\ L_{+n} &\mapsto \tilde{L}_{+n} := \frac{1}{n}L_{+n} \end{aligned} \quad \longmapsto \left\{ \begin{array}{l} [\tilde{L}_{+n}, \tilde{L}_{-n}] = 2\tilde{L}_0 \\ [\tilde{L}_{\pm n}, \tilde{L}_0] = \pm \tilde{L}_{\pm n} \end{array} \right\} sl(2, \mathbb{C}). \quad (3)$$

The corresponding finite transformations are of the form:

$$g_n(z) := \left(\frac{\alpha z^n + \beta}{\bar{\beta} z^n + \bar{\alpha}} \right)^{\frac{1}{n}}, \quad \begin{pmatrix} \alpha & \beta \\ \bar{\beta} & \bar{\alpha} \end{pmatrix} \in PSU(1, 1). \quad (4)$$

They leave the unit-circle S^1 invariant.

One may equally well represent the CFT_2 on a product of lines, $\mathbb{R} \times \mathbb{R}$ -spacetime (the "non-compact-picture"). The coordinate transformation from the circle to the line is provided by the stereographic projection (Cayley-transformation):

$$S^1 \setminus \{-1\} \ni z \mapsto x(z) := -i \frac{z-1}{z+1} \in \mathbb{R}, \quad -1 \mapsto \infty. \quad (5)$$

The global symmetry group $PSU(1, 1)$ transforms in this process isomorphically into the real group $PSL(2, \mathbb{R})$ [7]:

$$\mathbb{R} \ni x \mapsto \hat{g}(x) := \frac{ax + b}{cx + d}, \quad ad - bc = 1.$$

It is slightly more cumbersome to handle the transformations analogue to (eq. 4) in the non-compact picture. For this reason we perform the calculations in the compact picture representation.

Schroer and Wiesbrock take as a paradigm of their discussion the $U(1)$ -current-algebra on the circle. The constituting relation of this $U(1)$ -algebra is the current-current commutation-relation (with the circle as base-space):

$$[J(z), J(w)] = -\partial_z \delta(z - w).$$

In order to bring the algebraic ansatz [16] to the stage one has to smear the currents by real testfunctions on the circle:

$$J(f) := \int_{S^1} \frac{dz}{2\pi i} f(z) J(z).$$

The bounded operators $e^{iJ(f)}$ give rise to a von Neumann algebra, i.e. weakly closed algebra of bounded operators:

$$\mathcal{W}(\mathcal{I}) := \left\{ W(f) := e^{iJ(f)} \Big|_{\text{supp}(f) \subseteq \mathcal{I} \subset S^1} \right\}'' ,$$

which is called (local) Weyl-algebra [8, 17]. The double prime indicates here the double-commutant which after a theorem by von Neumann (double-commutant-theorem [17]) equals the weak closure of the algebra generated by the $e^{iJ(f)}$. The net of Weyl-algebras $\{S^1 \supset \mathcal{I} \mapsto \mathcal{W}(\mathcal{I})\}$ fulfills the postulates of algebraic quantum field theory, in particular locality [10].

The one-parameter group of dilations in $PSU(1, 1)$ has the following form:

$$Dil(t)z = \frac{ch(\pi t)z + sh(\pi t)}{sh(\pi t)z + ch(\pi t)}, \quad t \in \mathbb{R}, z \in S^1. \quad (6)$$

These mappings have the points $\{1, -1\} \in S^1$ as fixpoints. The upper and lower semi-circle, S^1_+ respectively S^1_- , are mapped by dilations onto themselves. A dilation-group attached to an arbitrary, proper interval $\mathcal{I} \subset S^1$ (mapping this interval onto itself) is constructed as follows:

$$PSU(1, 1) \ni Dil_{\mathcal{I}}(t) := g_{\mathcal{I}}^{-1} Dil(t) g_{\mathcal{I}}, \quad g_{\mathcal{I}} \in PSU(1, 1), \quad g_{\mathcal{I}} \mathcal{I} = S^1_+. \quad (7)$$

The interval $\mathcal{I}$ is mapped bijectively to the upper semi-circle, dilated and mapped back as one can see in the following diagram:

$$\begin{array}{ccc} \mathcal{I} & \xrightarrow{Dil_{\mathcal{I}}} & \mathcal{I} \\ g_{\mathcal{I}} \downarrow & & \uparrow g_{\mathcal{I}}^{-1} \\ S^1_+ & \xrightarrow{Dil} & S^1_+ \end{array}$$

The representation of the one-parameter group of dilations (eq. 7) gives (a geometric realization of) the unique modular group $\{\Delta_{\mathcal{I}}^{it}; t \in \mathbb{R}\}$ [16, 17] of the (vacuum) tuple $(\mathcal{W}(\mathcal{I}), \omega_0)$. For the case of the upper and lower semi-circle which in the non-compact picture become positive and negative lightrays this follows from the work of Bisognano and Wichmann [4]. It is a peculiarity of CFT_2 that arbitrary intervals can be mapped onto the upper (or lower) semi-circle which therefore allows to identify the modular group of algebras localized in these regions with the above constructed dilations. The vacuum expectation values of Weyl-operators obey the *KMS*-condition

[11, 16, 17] with respect to (the representation of) the one-parameter group of dilations [12]:

$$\omega_0(W(f) \text{Ad}[U_{\text{Dil}_{\mathcal{I}}(t)}](W(g))) \stackrel{\text{KMS}}{=} \omega_0(\text{Ad}[U_{\text{Dil}_{\mathcal{I}}(t+i)}](W(g)) W(f)) \quad (8)$$

$W(f), W(g) \in \mathcal{W}(\mathcal{I})$. In the case of the vacuum state this is a necessary and sufficient condition to identify uniquely the one-parameter group of dilations as the modular group [18] mentioned above.

By a simple reparametrization of the unit-circle $S^1 \subset \mathbb{C}$ in terms of the conformal mapping $S^1 \ni z \mapsto z^n$, $1 < n \in \mathbb{N}$, Schroer and Wiesbrock construct a geometrical state ω_2 for the case $n = 2$. We shall henceforth refer to it as F-S state. This state is shown to be invariant under transformations of the form (eq. 4) for $n = 2$:

$$g_2(z) := \left(\frac{\alpha z^2 + \beta}{\bar{\beta} z^2 + \bar{\alpha}} \right)^{\frac{1}{2}}, \quad \begin{pmatrix} \alpha & \beta \\ \bar{\beta} & \bar{\alpha} \end{pmatrix} \in PSU(1, 1). \quad (9)$$

For intervals:

$$\mathcal{I}^{\frac{1}{2}} := \mathcal{I}_1 \cup \mathcal{I}_2, \quad \mathcal{I}_i \xrightarrow{z^2} \mathcal{I}, \quad i = 1, 2, \quad (10)$$

the modified dilations act the following way:

$$\begin{aligned} \text{Dil}_{2,\mathcal{I}}(t)(\bullet) &:= (\text{Dil}_{\mathcal{I}}(t)(\bullet)^2)^{\frac{1}{2}} \\ &= (g_{\mathcal{I}}^{-1} \text{Dil}(t) g_{\mathcal{I}}(\bullet)^2)^{\frac{1}{2}} \end{aligned} \quad (11)$$

(see the following diagram)

$$\begin{array}{ccc} \mathcal{I}^{\frac{1}{2}} & \xrightarrow{\text{Dil}_{2,\mathcal{I}}} & \mathcal{I}^{\frac{1}{2}} \\ (\bullet)^2 \searrow & & \nearrow (\bullet)^{\frac{1}{2}} \\ \mathcal{I} & \xrightarrow{\text{Dil}_{\mathcal{I}}} & \mathcal{I} \\ g_{\mathcal{I}} \downarrow & & \uparrow g_{\mathcal{I}}^{-1} \\ S_+^1 & \xrightarrow{\text{Dil}} & S_+^1 \end{array}$$

The F-S state ω_2 may be defined in the following manner:

$$\omega_2(W(f) W(g)) := \omega_0(W(f_{\frac{1}{2}}) W(g_{\frac{1}{2}})), \quad W(f), W(g) \in \mathcal{W}(\tilde{\mathcal{I}}) \quad (12)$$

with $f_{\frac{1}{2}}(\bullet) := f((\bullet)^{\frac{1}{2}})$, $\text{supp}(f_{\frac{1}{2}}) \subset \mathcal{I}$, $\tilde{\mathcal{I}} \xrightarrow{z^2} \mathcal{I}$. It amounts to the following pointwise prescription for the current two-point-function:

$$\omega_2(J(z)J(w)) := 2z \, 2w \, \omega_0(J(z^2)J(w^2)). \quad (13)$$

At this point it is essential that the interval $\tilde{\mathcal{I}}$, i.e. localization region of the algebra $\mathcal{W}(\tilde{\mathcal{I}})$ in equation 12, does not contain opposite points $z, w \in$

S^1 , $\arg(z) - \arg(w) = 0, \text{ mod}(\frac{2\pi}{2})$. Otherwise the F-S state becomes non-faithful as one can see easily by the following example of an operator $W \in \mathcal{W}(\tilde{\mathcal{I}})$:

$$W := 1 - W(g)W(f) \quad (14)$$

$$f|_{\text{supp}(f)} = -g|_{\text{supp}(g)}, \quad \text{supp}(f) = -\text{supp}(g) \subset \tilde{\mathcal{I}}.$$

Localizing the algebra $\mathcal{W}$ in only one of the two intervals $\mathcal{I}_i \subset \mathcal{I}^{\frac{1}{2}}$, $i = 1, 2$, one is able to identify the modular group of the standard tuple $(\mathcal{W}(\mathcal{I}), \omega_2)$ ($\mathcal{I}$ stands for the chosen interval, i.e. $\mathcal{I}_1$ respectively $\mathcal{I}_2$) with the transformations in (eq. 11) by showing the *KMS*-property for the state ω_2 ⁴:

$$\omega_2(W(f) \text{Ad}[U_{\text{Dil}_{2,\mathcal{I}}(t)}](W(g))) \stackrel{\text{KMS}}{=} \omega_2(\text{Ad}[U_{\text{Dil}_{2,\mathcal{I}}(t+i)}](W(g)) W(f)). \quad (15)$$

Using equation 12 this may be reduced to the vacuum case:

$$\omega_0(W(f_{\frac{1}{2}}) \text{Ad}[U_{\text{Dil}_{\mathcal{I}}(t)}](W(g_{\frac{1}{2}}))) \stackrel{\text{KMS}}{=} \omega_0(\text{Ad}[U_{\text{Dil}_{\mathcal{I}}(t+i)}](W(g_{\frac{1}{2}})) W(f_{\frac{1}{2}})). \quad (16)$$

The faithful F-S state ω_2 on the algebra $\mathcal{W}(\mathcal{I})$ has a unique vector implementation $|\omega_2\rangle$ in the natural cone $\mathcal{P}_{|\omega_0\rangle}$ of the standard pair $(\mathcal{W}(\mathcal{I}), |\omega_0\rangle)$ [3, 16]:

$$\omega_2(W) = \langle \omega_2 | W | \omega_2 \rangle, \quad W \in \mathcal{W}(\mathcal{I}) \quad (17)$$

We want to point out in the following section that the above results of Schroer and Wiesbrock concerning the invariance of the modified states with respect to the modified $PSU(1, 1)$ group and the *KMS*-property of these states with respect to the modified dilations are general properties for any CFT_2 since the mentioned properties can -as we will show- be drawn through the substitutions $z \mapsto z^n$, $\forall n \in \mathbb{N}$.

III In the following we keep the pointwise prescription of fields. We generalize to arbitrary $n \in \mathbb{N}$, defining states $|\omega_n\rangle$ for local chiral primary fields $\phi(z)$ on S^1 by the identity:

$$\langle \omega_n | \prod_{k=1}^l \phi_k(z_k) | \omega_n \rangle := \prod_{s=1}^l (n z_s^{n-1})^{\Delta_s} \langle \omega_0 | \prod_{k=1}^l \phi_k(z_k^n) | \omega_0 \rangle. \quad (18)$$

Using the Möbius-invariance of the vacuum $|\omega_0\rangle$ one can show the invariance of $|\omega_n\rangle$ under transformations of the form (eq. 4) as follows:

$$\begin{aligned} \phi_i(z_i) &\longrightarrow (\partial_z \{g_n(z)\}|_{z=z_i})^{\Delta_i} \phi_i(g_n(z_i)) \\ \langle \omega_n | \prod_{k=1}^l \phi_k(z_k) | \omega_n \rangle &\longrightarrow \prod_{i=1}^l (\partial_z \{g_n(z)\}|_{z=z_i})^{\Delta_i} \langle \omega_n | \prod_{j=1}^l \phi_j(g_n(z_j)) | \omega_n \rangle \end{aligned}$$

⁴ Schroer and Wiesbrock did essentially the calculations in [1] whereas in [3] the correction, i.e. the limitation to only one of the two intervals as localization region was mentioned.

$$\begin{aligned}
& \stackrel{(eq.18)}{=} \prod_{i=1}^l (\partial_z \{g_n(z)\}|_{z=z_i})^{\Delta_i} \prod_{j=1}^l (n\{g_n(z_j)\}^{n-1})^{\Delta_j} \langle \omega_0 | \prod_{k=1}^l \phi_k(\{g_n(z_k)\}^n) | \omega_0 \rangle \\
& \stackrel{(eq.4)}{=} \prod_{i=1}^l (\partial_z \{g_n(z)\}|_{z=z_i})^{\Delta_i} \prod_{j=1}^l (n\{g_n(z_j)\}^{n-1})^{\Delta_j} \langle \omega_0 | \prod_{k=1}^l \phi_k(g(z_k^n)) | \omega_0 \rangle, \\
& \qquad \qquad \qquad g \in PSU(1,1) \\
& = \prod_{i=1}^l (\partial_z \{g_n(z)\}|_{z=z_i})^{\Delta_i} \prod_{j=1}^l (n\{g_n(z_j)\}^{n-1})^{\Delta_j} \times \\
& \qquad \qquad \qquad \prod_{r=1}^l \frac{1}{(\partial_z \{g(z)\}|_{z=z_r^n})^{\Delta_r}} \langle \omega_0 | \prod_{k=1}^l \phi_k(z_k^n) | \omega_0 \rangle \quad (19) \\
& \stackrel{(eq.18)}{=} \langle \omega_n | \prod_{i=1}^l \phi_i(z_i) | \omega_n \rangle.
\end{aligned}$$

For the one-parameter group of modified dilations:

$$\begin{aligned}
Dil_n(t)z &:= (Dil(t)z^n)^{\frac{1}{n}} \\
&= \left(\frac{ch(\pi t)z^n + sh(\pi t)}{sh(\pi t)z^n + ch(\pi t)} \right)^{\frac{1}{n}} \quad (20)
\end{aligned}$$

one would like the following *KMS*-relation to be satisfied:

$$\begin{aligned}
& (\partial_v \{Dil_n(t)v\}|_{v=z_l})^{\Delta_l} \langle \omega_n | \prod_{k=1}^{l-1} \phi_k(z_k) \phi_l(Dil_n(t)z_l) | \omega_n \rangle \\
& \stackrel{(eq.18)}{=} \prod_{i=1}^{l-1} (nz_i^{n-1})^{\Delta_i} [(\partial_v \{Dil_n(t)v\}|_{v=z_l})^{\Delta_l} (n\{Dil_n(t)z_l\}^{n-1})^{\Delta_l}] \times \\
& \qquad \qquad \qquad \langle \omega_0 | \prod_{k=1}^{l-1} \phi_k(z_k^n) \phi_l(Dil(t)z_l^n) | \omega_0 \rangle \\
& = \prod_{i=1}^l (nz_i^{n-1})^{\Delta_i} (\partial_v \{Dil(t)v\}|_{v=z_l^n})^{\Delta_l} \langle \omega_0 | \prod_{k=1}^{l-1} \phi_k(z_k^n) \phi_l(Dil(t)z_l^n) | \omega_0 \rangle \\
& \stackrel{!}{=} \prod_{i=1}^l (nz_i^{n-1})^{\Delta_i} (\partial_v \{Dil(t+i)v\}|_{v=z_l^n})^{\Delta_l} \times \\
& \qquad \qquad \qquad \langle \omega_0 | \phi_l(Dil(t+i)z_l^n) \prod_{k=1}^{l-1} \phi_k(z_k^n) | \omega_0 \rangle \\
& \stackrel{(eq.18)}{=} (\partial_v \{Dil_n(t+i)v\}|_{v=z_l})^{\Delta_l} \langle \omega_n | \phi_l(Dil_n(t+i)z_l) \prod_{k=1}^{l-1} \phi_k(z_k) | \omega_n \rangle. \quad (21)
\end{aligned}$$

We have managed here to prove the *KMS*-condition for the chiral part of the correlation function in the modified ($|\omega_0\rangle \mapsto |\omega_n\rangle$) theory provided that the *KMS*-condition holds in the unmodified theory. But the later is not precisely true. One picks in general a monodromy shift in the analytic continuation $t \mapsto t + is$, $s \in [0, 1]$ ⁵, which amounts to a full circle in complex space -returning to the same point on a different Riemann sheet. We may assume that the monodromy under consideration is diagonalized in a suitably chosen basis of conformal blocks and that the ensuing phase factors are compensated by the inverse phase factors of the anti-chiral block function (we assume the $\phi_{(i,j)}(z, \bar{z})$ to be scalar operators). The general situation is, what concerns the cancellation of phasefactors, faithfully represented by the simplified situation in the case of two-point-functions. The (vacuum-) 2-point-functions are already defined by conformal invariance (and locality: $\bar{\Delta} = \Delta$) to be of the form [15]:

$$\langle \omega_0 | \phi_1(z, \bar{z}) \phi_2(w, \bar{w}) | \omega_0 \rangle = \frac{C_{12}}{|z - w|^{4\Delta}}. \quad (22)$$

This 2-point-function fulfills the *KMS*-condition with respect to the one-parameter group of dilations $Dil(t) \in PSU(1, 1)$ (eq. 6) as one can show by direct calculation. In the case of arbitrary n -point-functions of local conformal fields $\phi_{(i,j)}(z, \bar{z})$ the above argument which holds for a rational conformal field theory [15] reduces the analytic structure essentially to the $U(1)$ -vertex-form, i.e. equation 22. Since the *KMS*-property holds for (eq. 22) it follows for the n -point-functions as well. Here it is important that one needs both, the chiral and anti-chiral part.

One has the following *KMS*-condition for general local conformal fields:

$$\begin{aligned} & (\partial_v \{Dil(t)v\}|_{v=z_l})^{\Delta_{i_l}} (\partial_{\bar{u}} \{Dil(-t)\bar{u}\}|_{\bar{u}=\bar{z}_l})^{\Delta_{j_l}} \times \\ & \langle \omega_0 | \prod_{k=1}^{l-1} \phi_{(i_k, j_k)}(z_k, \bar{z}_k) \phi_{(i_l, j_l)}(Dil(t)z_l, Dil(-t)\bar{z}_l) | \omega_0 \rangle \\ & \stackrel{KMS}{=} (\partial_v \{Dil(t+i)v\}|_{v=z_l})^{\Delta_{i_l}} (\partial_{\bar{u}} \{Dil(-t-i)\bar{u}\}|_{\bar{u}=\bar{z}_l})^{\Delta_{j_l}} \times \\ & \langle \omega_0 | \phi_{(i_l, j_l)}(Dil(t+i)z_l, Dil(-t-i)\bar{z}_l) \prod_{k=1}^{l-1} \phi_{(i_k, j_k)}(z_k, \bar{z}_k) | \omega_0 \rangle. \end{aligned} \quad (23)$$

A sufficient condition for locality is $\Delta_{i_p} = \Delta_{j_p}$, $p = 1, \dots, l$.

Going now to the Schroer-Wiesbrock ansatz, one has the following picture on the chiral and anti-chiral sector, respectively, and therefore the modified

⁵ In statistical physics one has the interval $[0, \beta]$, β the inverse temperature, and the states are called β -*KMS*-states [16, 17].

states $|\omega_{n,n}\rangle$:

$$\begin{aligned} \langle \omega_{n,n} | \prod_{k=1}^l \phi_{(i_k, j_k)}(z_k, \bar{z}_k) | \omega_{n,n} \rangle := \\ \prod_{s=1}^l (nz_s^{n-1})^{\Delta_{i_s}} \prod_{q=1}^l (n\bar{z}_q^{n-1})^{\Delta_{j_q}} \langle \omega_0 | \prod_{k=1}^l \phi_{(i_k, j_k)}(z_k^n, \bar{z}_k^n) | \omega_0 \rangle. \end{aligned} \quad (24)$$

Again, the *KMS*-condition with respect to $Dil_n(t)z := (Dil(t)z^n)^{\frac{1}{n}}$ (eq. 20) on both sectors "goes through covariantly" as in equation (eq. 21).

An illustration of what had been said above in a non-trivial setting (the $U(1)$ -current-algebra is a quasifree theory [8]) is provided by the theory of non-abelian currents [15]. The central relation is the current-current commutation relation (Kac-Moody-algebra, with the circle as base space):

$$[J^a(z), J^b(w)] = if^{abc} J^c(z) \delta(z-w) - k \delta^{ab} \partial_z \delta(z-w). \quad (25)$$

This relation allows it to calculate the m -point correlation function recursively by using the $m-1$ - and $m-2$ -point-function:

$$\begin{aligned} \langle \omega_0 | \prod_{i=1}^m J^{a_i}(z_i) | \omega_0 \rangle &\stackrel{(eq.25)}{=} \sum_{j=2}^m \frac{k \delta^{a_1 a_j}}{(z_1 - z_j)^2} \langle \omega_0 | \prod_{\substack{k=2 \\ k \neq j}}^m J^{a_k}(z_k) | \omega_0 \rangle + \\ &\sum_{j=2}^m \frac{i f^{a_1 a_j d}}{(z_1 - z_j)} \langle \omega_0 | \prod_{k=2}^{j-1} J^{a_k}(z_k) J^d(z_j) \prod_{l=j+1}^m J^{a_l}(z_l) | \omega_0 \rangle. \end{aligned} \quad (26)$$

One can therefore verify the *KMS*-property inductively since the 2-point function has the *KMS*-property. For the $|\omega_n\rangle$ state one gets the following recurrence relation:

$$\begin{aligned} \langle \omega_n | \prod_{i=1}^m J^{a_i}(z_i) | \omega_n \rangle &\stackrel{(eq.18)}{=} \sum_{j=2}^m \frac{(nz_1^{n-1})(nz_j^{n-1}) k \delta^{a_1 a_j}}{(z_1^n - z_j^n)^2} \langle \omega_n | \prod_{\substack{k=2 \\ k \neq j}}^m J^{a_k}(z_k) | \omega_n \rangle + \\ &\sum_{j=2}^m \frac{(nz_1^{n-1}) i f^{a_1 a_j d}}{(z_1^n - z_j^n)} \langle \omega_n | \prod_{k=2}^{j-1} J^{a_k}(z_k) J^d(z_j) \prod_{l=j+1}^m J^{a_l}(z_l) | \omega_n \rangle. \end{aligned}$$

The *KMS*-condition with respect to $Dil_n(t)z := (Dil(t)z^n)^{\frac{1}{n}}$ (eq. 20) can be shown inductively as well, since the substitution $z \mapsto z^n$ goes through covariantly and therefore the *KMS*-property for the vacuum carries over to the new state.

For the identification with the modular group mentioned above one needs von Neumann algebras. It is a subtle problem to proceed from smeared unbounded field operators to local algebras of bounded operators. By using

bounded functions of unbounded local operators as in the case of the $U(1)$ -current-algebra one has to ensure locality which might be not conserved by this mapping. Bisognano and Wichmann gave certain sufficient conditions to identify the von Neumann algebras to which the (Wightman-) field-algebras are affiliated [4, 17]. Generally it is a non-trivial task to verify these conditions. In the case of non-abelian currents, this is possible and one can therefore speak of von Neumann algebras, respectively the associated modular group.

IV We have seen that the localization region of the algebras has to exclude opposite points to guarantee the faithfulness of the F-S state ω_n . In the following we will stick to the case $n = 2$ for simplicity. Since both intervals $\mathcal{I}_{1,2}$ (eq. 10) are allowed for localizing the algebra it seems natural to have a closer look to the two-interval, i.e. multilocalized, algebra $\mathcal{W}(\mathcal{I}_1) \vee \mathcal{W}(\mathcal{I}_2) = \mathcal{W}(\mathcal{I}_1 \cup \mathcal{I}_2) =: \mathcal{W}(\mathcal{I}^{\frac{1}{2}})$. The demand for faithfulness prohibits the direct use of the F-S state ω_2 . There is however a way to enforce faithfulness by applying the split-property [16, 9]⁶.

For two spacelike regions, i.e. disjoint intervals, the split-property implies the following:

$$\mathcal{W}(\mathcal{I}_1) \vee \mathcal{W}(\mathcal{I}_2) \simeq \mathcal{W}(\mathcal{I}_1) \otimes \mathcal{W}(\mathcal{I}_2). \quad (27)$$

This reflects in a certain sense the statistical independence of the algebras. The state ω_2 , faithful and normal over either interval $\mathcal{I}_{1,2}$ can be extended to a product state, faithful and normal over $\mathcal{W}(\mathcal{I}^{\frac{1}{2}})$, i.e.:

$$\omega_2^p(WV) = \omega_2(W)\omega_2(V), \quad W \in \mathcal{W}(\mathcal{I}_1), \quad V \in \mathcal{W}(\mathcal{I}_2) \quad (28)$$

The modular group also splits and one can show the following theorem:

Theorem: The modular group $\sigma_{\omega_2^p}^t$ of the faithful product state ω_2^p on the algebra $\mathcal{W}(\mathcal{I}^{\frac{1}{2}})$ is given by the geometric action of $Dil_2(t)$. Moreover the unitary implementer $U_{Dil_2(t)}$ whose infinitesimal generator is a linear combination of $L_{\pm 2}$

$$\sigma_{\omega_2^p}^t = Ad[U_{Dil_2(t)}] \quad (29)$$

is the $\Delta_{\omega_2^p}^{it}$ modular object of the state vector $|\eta_2\rangle$ which represents the faithful product state ω_2^p in the positive cone of $(\mathcal{W}(\mathcal{I}^{\frac{1}{2}}), |\omega_0\rangle)$.

Since the arguments are entirely similar to those which demonstrated that $U_{Dil_2(t)}$ was the Δ^{it} modular group of the faithful state ω_2 on either $\mathcal{W}(\mathcal{I}_1)$ or $\mathcal{W}(\mathcal{I}_2)$, we omit the details. The crucial point is the invariance:

$$\omega_2^p(Ad[U_{Dil_2(t)}](WV)) = \omega_2^p(Ad[U_{Dil_2(t)}](W) Ad[U_{Dil_2(t)}](V))$$

⁶Another way might be the use of a (minimal) projector E implying the faithfulness of ω_2 on the reduced algebra EWE .

$$\begin{aligned}
&= \omega_2(Ad[U_{Dil_2(t)}](W)) \omega_2(Ad[U_{Dil_2(t)}](V)) \\
&= \omega_2(W) \omega_2(V) \\
&= \omega_2^p(WV)
\end{aligned} \tag{30}$$

which uses the previously established invariance of ω_2 . The aforementioned lack of faithfulness of the state ω_2 on the multi-interval algebra $\mathcal{W}(\mathcal{I}^{\frac{1}{2}})$ is related to the geometric nature of the double interval modular group $\sigma_{\omega_2^p}^t$.

The remarkable fact that the pairs $(\mathcal{W}(\mathcal{I}_i), \omega_2)$, $i = 1, 2$, $(\mathcal{W}(\mathcal{I}^{\frac{1}{2}}), \omega_2 \times \omega_2)$ share the same modular group action and (in the appropriate positive cones) have the same implementing modular unitaries is also related to this. The factorization of the ω_0 vacuum would not lead to such a situation.

This result raises the question whether both modular objects of the double interval situation $(\mathcal{W}(\mathcal{I}^{\frac{1}{2}}), |\eta_2\rangle)$ can be geometric. Formally the candidate for a geometric J is the "rotated" TCP transformation $z \rightarrow -\bar{z}$ which maps for instance the first quarter-circle $S_1^1 := [0, \frac{\pi}{2}]$ to the second quarter-circle $S_2^1 := [\frac{\pi}{2}, \pi]$ and the third quarter-circle S_3^1 to the fourth S_4^1 ⁷ (this transformation has the same geometric effect as the analytically continued $\Delta_{\omega_2^p}^{\frac{1}{2}}$). However, the replacement of Haag duality by an inclusion:

$$\mathcal{W}((\mathcal{I}_1 \cup \mathcal{I}_2)') \subset \mathcal{W}(\mathcal{I}_1 \cup \mathcal{I}_2)' \tag{31}$$

and its explanation in term of superselection sectors shows that the true modular involution has in addition to the geometric part an algebraic modification. A calculation of J for abelian current models seems to be feasible. Via the inverse algebraic lightfront holography [3] these geometric chiral transformations correspond to fuzzy symmetries in the original net.

V It was shown that Dil_n (having $2n$ fixpoints instead of two as Dil) is the modular group of the standard tuple $(\mathcal{W}(\mathcal{I}), \omega_n)$. The interval $\mathcal{I}$ is forbidden to contain points $z, w \in S^1$, $arg(z) - arg(w) = 0, \text{ mod}(\frac{2\pi}{n})$.

With regard to the modular origin of the Witt-Virasoro basis it is sufficient to extend the construction up to the $n = 2$ case due to the relation (1).

In the case of multi-interval, i.e. multilocalized Weyl-algebras (eq. 27), the transformations (eq. 29) can be identified with the modular group of the tuple $(\mathcal{W}(\mathcal{I}^{\frac{1}{2}}), |\eta_2\rangle)$. Going to higher n by using the split-property is not totally straightforward.

Following the program of algebraic quantum field theory this result underlines the special rôle played by modular theory in general local quantum field theory. Schroer and Wiesbrock and Schroer and Fassarella [3] suggest using modular theory as a tool to explore "fuzzy" symmetries i.e. symmetries which do not originate from the classical Noether setting. They

⁷The quarter-circle algebras $\mathcal{W}(S_{1,3}^1)$ related to the algebra localized on the upper semi-circle $\mathcal{W}(S_+^1)$ provide a natural example.

also propose to investigate the relation of their findings with the notion of *half-sided-modular-inclusion(intersections)* [2, 5, 13]. See [6, 14] for a recent account and further references to the results in [2, 5, 13].

To investigate the latter was out of scope of the present work. Here, we showed that a new ansatz of Schroer and Wiesbrock (corrected in [3]) can be dealt with from a more general point of view, since the *KMS*-property of the vacuum state (a general property for local fields in the vacuum-setting) carries over covariantly to the set of new F-S states. The argument for the general validity of the *KMS*-property given above is true for rational conformal field theories.

Multilocalized algebras can be defined by using the split-property which implies the faithful states to be a product of the new constructed F-S states. The modular group naturally splits into a product of two copies of the modified dilations whereas the modular conjugation needs some further investigation.

Note to be added: We are indebted to Prof. Schroer for pointing out to us the incorrectness of the implicitly assumed faithfulness of the new states with respect to multilocalized algebras in [1]. Section 4 is based on notes by Prof. Schroer.

Acknowledgements: The author is grateful to Prof. Dr. R. Flume for suggesting this work. Also, we would like to thank Kayhan Ülker and Nicolai von Rummell for help and wise comments when preparing this work. I thank also H. Gutmann for a careful reading of the manuscript.

References

- [1] B. Schroer, H.-W. Wiesbrock, "Modular Theory and Geometry", *Rev. Math. Phys.* **12** (2000) 139.
- [2] H.-W. Wiesbrock, "Conformal quantum field theory and half-sided modular inclusions of von-Neumann algebras", *Commun. Math. Phys.* **158** (1993) 537
- [3] B. Schroer, L. Fassarella, "The Fuzzy Analog of Chiral Diffeomorphisms in higher dimensional Quantum Field Theories", preprint hep-th/0106064.
- [4] J. Bisognano, E. Wichmann, "On the duality condition for a hermitian scalar field", *J. Math. Phys.* **16** (1975) 985.
- [5] H.-J. Borchers, "On the use of modular groups in quantum field theory", *Ann. Inst. Henri Poincaré* **63** n°4(1995) 331

- [6] D. Buchholz, "Algebraic Quantum Field Theory: A Status Report" preprint math-ph/0011044
- [7] F. Gabbiani, J. Fröhlich, "Operator Algebras and Conformal Field Theory", App.I, *Commun. Math. Phys.* **155** (1993) 569
- [8] D. Buchholz, G. Mack, I. Todorov, "The current algebra on the circle as a germ of local field theories", *Nucl. Phys.***B**, Proc. Suppl. **56** (1988) 20
- [9] B. Schroer, "Particle Physics and quantum field theory at the turn of the century: Old principles with new concepts", *J. Math. Phys.***45** (2000) 3801.
- [10] D. Buchholz, H. Schulz-Mirbach, "Haag Duality in conformal quantum field theory", *Rev. Math. Phys.* **2** (1990) 105
- [11] R. Haag, N. Hugenholtz, M. Winnink, "On the equilibrium state in quantum statistical mechanics", *Commun. Math. Phys.* **5** (1967) 215
- [12] J. Yngvason, "A note on essential duality", *Lett. Math. Phys.***31** (1995) 127
- [13] H.-W. Wiesbrock, "Symmetries and half-sided modular inclusions of von-Neumann algebras", *Lett. Math. Phys.***29** (1993) 107
- [14] R. Kähler, H.-W. Wiesbrock, "Modular Theory and the reconstruction of four-dimensional quantum field theory ", *J. Math. Phys.***42** (2001) 74.
- [15] P. Di Francesco, P. Mathieu, D. Sénéchal, "Conformal Field Theory", Springer Verlag (Heidelberg 1997)
- [16] R. Haag, "Local Quantum Physics", 2nd ed., Springer Verlag (Berlin 1996)
- [17] O. Bratelli, D. W. Robinson, "Operator Algebras and Quantum Statistical Mechanics", Vol.1, 2, 2nd ed., TMP Springer Verlag (Berlin 1996)
- [18] R. V. Kadison, J. R. Ringrose, "Fundamentals of the Theory of Operator Algebras II", Academic Press (New York 1986)